



2025 Cyanobacteria Monitoring Report for Town of Harwich, Harwich, Massachusetts

December 17, 2025

Prepared for the town of Harwich
P.O. Box 207, Harwichport, MA 02646
Contact: Stephanie Ridenour at stephanie.ridenour@harwich-ma.gov
Amy Usowski ausowski@town.harwich.ma.us

Prepared by the Association to Preserve Cape Cod
482 Main Street, Dennis, MA 02638
Phone: (508) 619-3185
apcc.org
Contact: Dr. Julie Hambrook at jhambrook@apcc.org

Table of Contents

1. SUMMARY	4
2. BACKGROUND	5
3. METHODS AND MONITORING OVERVIEW	6
4. ASSESSING RISK	7
5. RESULTS.....	9
5.1 Cyanobacteria Presence and Biomass.....	9
5.2 Toxin Screening and Confirmatory Testing	19
5.3 Advisory Issuance and Resolution	20
6. RECOMMENDATIONS	21
7. ACKNOWLEDGEMENTS.....	23
8. REFERENCES	23
9. APPENDIX 1-4.....	26
Appendix 1. Resources on Cyanobacteria	26
Appendix 2. APCC's Cyanobacteria Risk Categories.....	28
Appendix 3. Bucks Pond, Cornelius Pond, Hawksnest Pond, Hinkley Pond, Josephs Pond, Sand Lake, Sand Pond, Skinequit Pond, and West Reservoir 2021-2025 Cyanobacteria Risk Comparison.....	31
Appendix 4. Sampling Locations.....	34

Tables

Table 1. Summary of all sampling events that took place in Harwich for the town of Harwich in 2025. ...	4
Table 2. Summary of all toxin testing that took place in Harwich for the town of Harwich in 2025 and of the number of measurements that exceeded MDPH guidelines per sampling location.	4
Table 3. Microcystin samples collected at each sampling location during 2025.	19
Table 4. Anatoxin-a samples collected at each sampling location during 2025.....	20

Figures

Figure 1. APCC 2025 Cyanobacteria Risk Tiers.....	8
Figure 2. Sampling Locations for town of Harwich in 2025.	9
Figure 3. Town of Harwich: Median Bloom-Forming-Colony phycocyanin measurements per monitored pond in town of Harwich across the 2025 season.	10
Figure 4. Bucks Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event.....	11
Figure 5. Cornelius Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event.	12
Figure 6. Hawksnest Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event.	13
Figure 7. Hinkleys Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event.	14

Figure 8. Josephs Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event.	15
Figure 9. Sand Lake: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event.....	16
Figure 10. Sand Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event.....	17
Figure 11. Skinequit Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event.	18
Figure 12. West Reservoir: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event.	19
Figure 13. Town of Harwich summary of the risk tiers reflected on APCC's Cyano Map throughout the 2025 season.....	21

1. SUMMARY

In 2025, the Association to Preserve Cape Cod (APCC), with the help of the staff at Barnstable County Lab, continued cyanobacteria monitoring in Harwich for the town of Harwich following similar monitoring conducted in 2021 through 2024. View Table 1 for an overview of all sampling events conducted by APCC in 2025 from June 3, 2025, through November 4, 2025. For ponds being monitored for the Cape Cod Regional Pond Monitoring Program (CCRPMP) that were not being monitored for cyanobacteria - a grant from the Southeast New England Coastal Watershed Restoration Program to the Cape Cod Commission provided funding for cyanobacteria monitoring of Cornelius Pond, Hawksnest Pond, and Sand Lake during 2024 and 2025.

Table 1. Summary of all sampling events that took place in Harwich for the town of Harwich in 2025.

Pond Name & sampling location	Sampling Frequency	# of Sampling Events	Highest Risk Tier reached in 2025	Highest Town Action taken in 2025
Bucks	Biweekly	16	Potential for Concern	Warning
Cornelius	Monthly	3	Acceptable	-
Hawksnest	Monthly	2	Acceptable	-
Hinckleys	Biweekly	12	Acceptable	-
Josephs	Biweekly	12	Acceptable	-
Sand Lake	Monthly	3	Acceptable	-
Sand Pond	Biweekly	14	Potential for Concern	Warning
Skinequit	Biweekly	12	Acceptable	-
West Reservoir	Biweekly	20	Potential for Concern	Warning
Total	-	94	-	-

View Table 2 for an overview of all toxin testing conducted throughout the 2025 season.

Table 2. Summary of all toxin testing that took place in Harwich for the town of Harwich in 2025 and of the number of measurements that exceeded MDPH guidelines per sampling location.

Pond Name & sampling location	# of Microcystin Strip Tests (APCC)	# of Microcystin ELISA Tests (BCDHE)	# of Anatoxin-a Strip Tests (APCC)	# of Anatoxin-a ELISA Tests (BCDHE)	# of Microcystin exceedances (>8ug/L)	# of Anatoxin-a exceedances (>60ug/L)
Bucks	4	1	-	-	-	-
Skinequit	-	-	1	-	-	-
West Reservoir	9	1	4	2	-	-

APCC shared all monitoring results with the town of Harwich, and the public throughout the season via emailed updates, e-newsletters, frequent updates to our online map at <https://apcc.org/our-work/science/community-science/cyanobacteria/>, and written reports,

including this report. This report should be printed in color, as some sections are color-coded. For the full dataset see Attachment 1.

2. BACKGROUND

APCC's Cyanobacteria Monitoring Program partners with officials at the municipal, county, state, and federal levels, as well as local pond associations and residents, to conduct cyanobacteria monitoring in select Cape Cod ponds. Each season, water samples are collected biweekly for monitoring of cyanobacteria and the results are shared with local officials and the general public through reports, emails, and an interactive map of monitoring results provided on our website (<https://apcc.org/cyano>). APCC's goals are to raise public awareness of the health and ecological risks posed by harmful cyanobacteria blooms (HCBs), collect scientific monitoring data, help inform proper responses to cyanobacteria blooms, protect public health, monitor priority ponds across the Cape, and motivate public action to address the causes of HCBs by improving water quality.

Cyanobacteria are an ancient group of photosynthetic microorganisms common in freshwater systems on Cape Cod, in the U.S., and worldwide. Under the right conditions, they can multiply rapidly and form harmful cyanobacteria blooms (HCBs). According to the Centers for Disease Control and Prevention (CDC), certain common cyanobacteria genera can produce toxins known as cyanotoxins that can be harmful to humans (https://www.cdc.gov/harmful-algal-blooms/about/?CDC_AAref_Val=https://www.cdc.gov/habs/general.html). There are many different genera of cyanobacteria present in Cape Cod freshwater ponds, with *Microcystis* and *Dolichospermum* being two of the most common. Cyanobacteria can produce a variety of toxins, many of which are still not fully understood. Our monitoring efforts have focused primarily on one of the best studied toxins, microcystin, which is commonly produced by *Microcystis*. As of 2025, APCC has also begun more regular monitoring for anatoxin-a, a toxin commonly produced by *Dolichospermum* as well as other cyanobacterial genera. HCBs have increased worldwide, including in the U.S., due to nutrient enrichment and rising water temperatures due to climate change. As the occurrence of HCBs increases the need for cyanobacteria monitoring and awareness of potential health risks has increased. Additional resources on cyanobacteria are provided in Appendix 1.

Cape Cod ponds are commonly used for recreation, including swimming, boating, paddle boarding, fishing, as well as for dog walking and swimming. Due to the increasing prevalence of HCBs and the resulting increased threat of public exposure to cyanobacteria and their toxins, MDPH has toxin standards for microcystin and anatoxin-a and provides guidelines for municipal officials to post and remove advisories at ponds based on established thresholds for cyanobacteria risks in recreational waters (<https://www.mass.gov/info-details/guidelines-for-cyanobacteria-at-recreational-freshwater-locations>). Frequent cyanobacteria monitoring of ponds provides fact-based data for resource managers to track cyanobacteria trends in their ponds throughout the season, apply relevant public health criteria, and proactively post and remove recreational advisories. Cyanobacteria monitoring data also provides information on pond health and water quality and helps to address data gaps caused by lack of conventional pond water quality monitoring data.

3. METHODS AND MONITORING OVERVIEW

APCC and Barnstable County Lab staff conducted cyanobacteria monitoring at nine locations in Harwich during the 2025 season, following methods outlined in APCC's Quality Assurance Project Plan (QAPP), based on the EPA's Cyanobacteria Monitoring Collaborative (CMC) QAPP. These methods have been in use since the program's inception in 2017 and ensure that monitoring is scientifically sound and consistent across seasons.

Field sampling occurred primarily on a biweekly basis from June 3, 2025, to November 4, 2025, with weekly monitoring triggered when a pond reached the "Potential for Concern" or "Use Restriction Warranted" risk categories.

A total of 94 sampling events were conducted, and collected samples were analyzed for cyanobacteria genus composition and phycocyanin, a pigment used as a proxy for cyanobacteria biomass. APCC and Barnstable County Lab staff collects several types of phycocyanin samples to assess different parts of the water column and bloom conditions. These include a <50 µm sample (to assess small-celled cyanobacteria), a Whole-Lake-Water (WLW) sample (to assess overall presence in the water column), and Bloom Forming Colony (BFC) sample collected using a plankton net towed along the water surface. The BFC phycocyanin sample results are used for risk assessment purposes. Genus dominance was defined as $\geq 70\%$ of colonies under microscopy. For details on the methodology see the link to [Cape Cod Cyanobacteria Monitoring Program \(CCCMP\) Quality Assurance Project Plan \(QAPP\) 2024-2029](https://apcc.org/our-work/science/community-science/cyanobacteria/) on APCC's website <https://apcc.org/our-work/science/community-science/cyanobacteria/>.

Each sampling event also included visual scum assessments, which were documented using APCC's standardized photo guide (Attachment 2.) to classify confirmed cyanobacteria scums as "insignificant" or "significant." Confirmed significant scums are treated as potential toxin risk even in the absence of elevated phycocyanin levels.

Since 2022, APCC has partnered with the Barnstable County Department of Health and Environment (BCDHE) Water Quality Lab for testing of the cyanobacteria toxin microcystin. The County Lab is a state-certified facility that uses an EPA-approved ELISA protocol. In 2025, toxin testing continued under this partnership, now also including anatoxin-a testing.

To provide more timely toxin information in 2025 in advance of receiving results from BCDHE, APCC utilized a handheld AbraScan digital strip test reader with Gold Standard Diagnostics microcystin and anatoxin-a test strips as a screening tool. Screening was only conducted for samples from ponds categorized as "Potential for Concern" or "Use Restriction Warranted" based on scum and/or phycocyanin levels, and only when microscopic analysis identified dominance of *Microcystis* spp. (for microcystin testing) or *Dolichospermum* spp. (for anatoxin-a testing). Microcystin strip test results ≥ 4 µg/L were sent to the BCDHE Water Quality Lab for confirmatory analysis. Samples dominated by *Dolichospermum* spp. were generally not tested for microcystin, based on historic APCC data indicating a lack of correlation between this genus and microcystin production. Most anatoxin-a strip test detections were also forwarded to the BCDHE Water Quality Lab for confirmatory analysis, as the strip test screening method is still under evaluation.

Sampling results are provided in biweekly reports to local municipal officials and pond associations and uploaded to APCC's interactive cyanobacteria risk map (apcc.org/cyano), which updates every six hours on weekdays during the monitoring season and is publicly accessible. APCC also provides an email registry signup on our website designed to update interested residents and visitors when harmful cyanobacteria blooms are identified. A quick link to sign up for the Cyanobacteria Alert is on our homepage (<https://apcc.org/our-work/science/community-science/cyanobacteria/cyanobacteria-alert/>).

4. ASSESSING RISK

In 2025, APCC revised its cyanobacteria risk tier system to better reflect the conditions observed in Cape Cod ponds and to improve the consistency and clarity of public risk communication. These revisions were informed by analysis of APCC's cyanobacteria monitoring data (2022–2024), including phycocyanin levels and microcystin concentrations, as well as practical experience managing the program and coordinating with both MDPH and town health departments. The result is a system that offers standardized, science-based guidance to help communities assess potential health risks while leaving decisions on the issuance of Warnings and Public Health Advisories to local health officials.

The risk tier framework continues to use three categories; each tier is defined as pictured in the 2025 Risk Tier Table in Figure 1 and detailed description in Appendix 2.

APCC 2025 Cyanobacteria Risk Tiers

Acceptable

- A cyanobacteria scum was not detected, and the Bloom Forming Colony sample had a phycocyanin measurement <500ug/L.
- A cyanobacteria scum was detected but was determined to be visually insignificant and the Bloom Forming Colony sample had a phycocyanin measurement <100ug/L.

Potential for Concern

- A cyanobacteria scum was not detected but the Bloom Forming Colony sample had a phycocyanin measurement ≥500ug/L.
- A cyanobacteria scum was detected and determined to be visually significant and/or the Bloom Forming Colony sample had a phycocyanin measurement ≥100ug/L.

Stripes are added on the map if the town posts a warning that is not an official Public Health Advisory.*

Use Restriction Warranted

- The town posts a Public Health Advisory.
- Microcystin test measures ≥8 ppb (MassDPH guidance).
- Once a pond is categorized as Use Restriction Warranted it will remain in this category for two consecutive Acceptable sampling events (MassDPH guidance).

Stripes are added on the map if the town posts a Public Health Advisory and are removed once the town removes the Public Health Advisory. **If and when the town informs APCC of their action*

Figure 1. APCC 2025 Cyanobacteria Risk Tiers

While APCC does not rely on cyanobacteria cell counts, its risk categories are generally consistent with Massachusetts Department of Public Health (MDPH) guidelines, which recommend issuing advisories when cell counts exceed 70,000 cells/mL, visible scums are present, microcystin concentrations exceed 8 µg/L, or anatoxin-a concentrations exceed 60 µg/L.

APCC's risk map reflects through the use of stripes to show when a managing agency has issued an official Warning or Advisory. These are removed once the agency lifts the warning.

It's important to note that APCC does not issue Public Health Advisories. Instead, APCC provides independent, science-based data interpretation and standardized risk assessment to support informed decision-making across Cape Cod at both the town and personal level. Managing agencies retain full authority to issue Public Health Advisories based on their own protocols. At the same time, individual users can use APCC's consistent, Cape-wide risk assessments to better understand cyanobacteria conditions and make personal decisions based on their own level of risk tolerance.

2025 Sampling Locations




Figure 2. Sampling Locations for town of Harwich in 2025.

5. RESULTS

During the 2025 monitoring season, APCC and Barnstable County Lab staff collected, and APCC analyzed samples from nine locations within the town of Harwich (see Table 1). Monitoring focused on assessing cyanobacteria presence, biomass (via phycocyanin), and microcystin and anatoxin-a toxin levels under the revised Risk Tier framework.

5.1 Cyanobacteria Presence and Biomass

Cyanobacteria monitoring results, Risk Categories, and risk communication are described in this section. For each pond, a graph is provided to describe results and risk categories for each sampling event. A complete table of results is provided in Attachment 1 containing all data collected for the town of Harwich in 2025. The full Risk Category criteria are included in Appendix 2. See Figure 3 for the median phycocyanin measurements per pond across 2025 sampling events.




Figure 3. Town of Harwich: Median Bloom-Forming-Colony phycocyanin measurements per monitored pond in town of Harwich across the 2025 season. The larger and more yellow/red the dot, the higher the median biomass.

Pond 1. Bucks Pond:

During the 2025 monitoring season, Bucks Pond experienced changes in cyanobacteria levels beginning with a “Potential for Concern” on June 16 which triggered weekly sampling. The cyanobacteria population reduced in the subsequent weeks, which placed it in APCC’s “Acceptable” Risk Category until November 7, 2025 (Figure 4. below).




Figure 4. Bucks Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event. The dominant genus abbreviations on this graph are *Microcystis* (MC) and mixed (mix).

Pond 2. Cornelius Pond:

During the 2025 monitoring season, Cornelius Pond contained no concerning cyanobacteria results at the time and place of each sampling event, keeping the pond in APCC's "Acceptable" category for the entire season (see Figure 5 below).




Figure 5. Cornelius Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event. The dominant genus abbreviations on this graph are *Microcystis* (MC) and *Dolichospermum* (DS).

Pond 3. Hawksnest Pond:

During the 2025 monitoring season, Hawksnest Pond contained no concerning cyanobacteria results at the time and place of each sampling event, keeping the pond in APCC's "Acceptable" category for the entire season (see Figure 6 below). On August 5 APCC suspended cyanobacteria monitoring in Nickerson State Park. APCC was notified by the state Department of Conservation and Recreation (DCR) that APCC's access to the park for the purposes of data collection is contingent upon APCC obtaining a DCR research permit. While APCC submitted an application on August 7 the application has not been approved.





Figure 6. Hawksnest Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event. The dominant genus abbreviation on this graph is none (-).

During the 2025 monitoring season, Hinckleys Pond contained no concerning cyanobacteria results at the time and place of each sampling event, keeping the pond in APCC’s “Acceptable” category for the entire season (see Figure 7 below).



Pond 5. Josephs Pond:

During the 2025 monitoring season, Josephs Pond contained no concerning cyanobacteria results at the time and place of each sampling event, keeping the pond in APCC's "Acceptable" category for the entire season (see Figure 8 below).




Figure 8. Josephs Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event. The dominant genus abbreviations on this graph are *Dolichospermum* (DS), *Microcystis* (MC), mixed (mix), and none (-).

Pond 6. Sand Lake:

During the 2025 monitoring season, Sand Lake contained no concerning cyanobacteria results at the time and place of each sampling event, keeping the pond in APCC's "Acceptable" category for the entire season (see Figure 9 below).




Figure 9. Sand Lake: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event. The dominant genus abbreviations on this graph are *Dolichospermum* (DS), *Microcystis* (MC), and none (-).

Pond 7. Sand Pond:

During the 2025 monitoring season, Sand Pond experienced changes in cyanobacteria levels late in the season with a “Potential for Concern” on September 9, which triggered weekly sampling. The cyanobacteria population reduced in the subsequent weeks, which placed it in APCC’s “Acceptable” Risk Category for the remainder of the season (see Figure 10 below).




Figure 10. Sand Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event. The dominant genus abbreviations on this graph are *Dolichospermum* (DS), *Microcystis* (MC), and none (-).

Pond 8. Skinequit Pond:

During the 2025 monitoring season, Skinequit Pond contained no concerning cyanobacteria results at the time and place of each sampling event, keeping the pond in APCC's "Acceptable" category for the entire season (see Figure 11 below).




Figure 11. Skinequit Pond: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event. The dominant genus abbreviations on this graph are *Woronichinia* (WO), *Microcystis* (MC), *Oscillatoria* (Osc), mixed (mix), none (-), and other.

Pond 9. West Reservoir:

During the 2025 monitoring season, West Reservoir experienced changes in cyanobacteria levels beginning with a “Potential for Concern” on July 1, which triggered weekly sampling for majority of the monitoring season. The cyanobacteria population reduced on October 14, which placed it in APCC’s “Acceptable” Risk Category for the remainder of the season (see Figure 12 below).




Figure 12. West Reservoir: BFC PC, dominant genus (>70%), risk tier reflected on APCC's map per sampling event. The dominant genus abbreviations on this graph are *Dolichospermum* (DS), *Microcystis* (MC), mixed (mix), and other.

5.2 Toxin Screening and Confirmatory Testing

APCC conducted thirteen microcystin and five anatoxin-a strip tests during the 2025 season based on cyanobacteria genus dominance. A total of four samples were submitted for confirmatory laboratory analysis at BCDHE following preliminary field results that met or exceeded screening thresholds or were used to study screening result reliability. For all toxin results see Table 3 and 4 below.

Table 3. Microcystin samples collected at each sampling location during 2025.

Pond name	Site	Date	Strip Test Result (ug/L)	BCDHE Result (ug/L)
Bucks	1	6/24/2025	nd	-
Bucks	1	7/1/2025	nd	-
Bucks	1	7/8/2025	nd	-
Bucks	1	7/15/2025	~5	0.45

West Reservoir	1	7/29/2025	~2.5	2.2
West Reservoir	1	8/12/2025	nd	-
West Reservoir	1	8/19/2025	nd	nd
West Reservoir	1	8/26/2025	nd	-
West Reservoir	1	9/2/2025	nd	-
West Reservoir	1	9/9/2025	nd	-
West Reservoir	1	9/17/2025	nd	-
West Reservoir	1	9/23/2025	nd	-
West Reservoir	1	9/30/2025	nd	-

Table 4. Anatoxin-a samples collected at each sampling location during 2025.

Pond name	Site	Date	Strip Test Result (ug/L)	BCDHE Result (ug/L)
Skinequit	1	7/15/2025	nd	-
West Reservoir	1	7/22/2025	nd	-
West Reservoir	1	7/29/2025	nd	-
West Reservoir	1	8/6/2025	nd	-
West Reservoir	1	8/19/2025	~.4	nd
West Reservoir	1	9/23/2025	-	0.26

5.3 Advisory Issuance and Resolution

Throughout the season, two ponds had official Warning or Advisory postings related to cyanobacteria issued by a managing agency. An overview of the Risk Tiers each monitored pond was assigned to during the 2025 season in Figure 13.




Figure 13. Town of Harwich summary of the risk tiers reflected on APCC's Cyano Map throughout the 2025 season. If town action was taken and a Warning or a Public Health Advisory was posted the “time window” has a bold black frame. If there is no black bold frame, no official town action was taken. If a time window for a given pond is blank, no sampling event took place during that window. If multiple sampling events took place during a given time window, the highest assigned risk tier during that time is reflected here.

6. RECOMMENDATIONS

Based on the results from the 2025 monitoring season and previous monitoring work, APCC provides the following recommendations:

Recommendation 1: Continue annual cyanobacteria monitoring. Monitoring over multiple years for full seasons would provide greater understanding of the cyanobacteria community in the ponds in the town of Harwich. More seasons of data will allow us to draw better predictions year after year. Continued monitoring will also allow for the ability to track degradation in the ponds as increased occurrence of harmful cyanobacteria blooms point to larger issues of pond impairment. Monitoring efforts will shed light on the ponds most in need of protection and restoration.

Recommendation 2: Reduce nutrient loading to freshwater ponds. Algal blooms and cyanobacteria blooms in ponds are associated with nutrient loading to ponds. Furthermore, there is mounting evidence that nitrogen, as well as phosphorus, is involved in stimulating algal blooms in freshwater ponds. Residents living adjacent to ponds and within the pond watershed should reduce sources of nutrient pollution flowing from their properties towards the pond. Excess fertilizer use, septic systems around ponds, poor stormwater management infrastructure, and a lack of adequate vegetation buffers exacerbate nutrient loading of ponds. Eliminating, or at

the very least minimizing, fertilizer use, replacing septic systems with municipal collection and treatment that remove more nutrients, treating stormwater runoff before it enters wetlands or ponds, and planting native vegetated buffers where none exist will help to protect pond water quality.

Recommendation 3: Recognize that scientific understanding of the causes of HCBs continues to evolve. In addition to managing nutrients, climate change, including the currently warming atmosphere and altered rainfall patterns, is believed to play a significant role in the increasing frequency and intensity of harmful cyanobacteria blooms (<https://www.publish.csiro.au/MF/MF18392>). Residents and officials should understand that there may be many factors that cause HCBs on Cape Cod. Continued monitoring of cyanobacteria and water quality will lead to increased understanding and awareness, a safer public, and hopefully improved health of our freshwater ponds.

Recommendation 4: Consider carefully before planning or undertaking pond restoration and protection options. If a pond is impacted by HCBs, here are some steps to consider:

- Identify the important uses and desirable features of the pond and surrounding areas.
- Consider the natural evolution of a pond over time, from open water to vegetated marsh to wet meadow.
- Identify potential causes of HCBs (e.g., stormwater runoff, fertilized lawns, etc.);
- Identify potential actions to promote pond health and reduce HCBs.
 - For a comprehensive list of actions that residents, municipalities, and state agencies can take to promote pond health, visit APCC's State of the Waters: Cape Cod website (<https://capecodwaters.org/action-plan/#ponds-hom>), specifically, the Action Plan for ponds.
 - Under the Freshwater Initiative, the Cape Cod Commission developed a Pond Restoration Technologies matrix to identify potential methods for pond protection and restoration (<https://ww2.capecodcommission.org/fwi/index.html>). Each pond is unique, and restoration technology that works for one pond may not work well for another. Evaluate whether potential actions to reduce HCBs may impact other pond characteristics or aquatic organisms that are important for pond health. Treating a single problem without considering the effects on other pond resources could potentially harm the entire pond ecosystem and its resources.
- Develop a comprehensive plan for pond protection and restoration that addresses the causes as well as the symptoms of impaired pond health.
- Planning for pond restoration should be done to address not only short-term solutions but also medium-term and long-term solutions.
- Develop a watershed management plan and continue to engage residents around the pond and within the watershed about best land care practices, pond ecology, etc.

7. ACKNOWLEDGEMENTS

APCC collaborates with many local, regional, state and federal partners, including organizations, homeowner associations, pond associations, water quality committees, municipal staff from Cape Cod, and state and federal agencies and organizations. Partners include scientists affiliated with Barnstable County Department of Health and the Environment, the Cape and Islands Health Agents Coalition, Massachusetts Department of Public Health, Massachusetts Department of Environmental Protection, the U.S. Environmental Protection Agency, and Massachusetts Bays National Estuary Partnership. Funding was provided by the Mary-Louise Eddy and Ruth N. Eddy Foundation, the Bilezikian Family Foundation, the Cape Cod 5 Foundation, the Eversource Energy Foundation, Inc., the Palmer Foundation, private foundation grants, dues, and donations from APCC members.

APCC wishes to thank the following individuals and organizations for their support of this project: Barry H. Rosen, Ph.D. of Florida Gulf Coast University for providing a taxonomic workshop and keys for identification of freshwater cyanobacteria and green algae, Mindy L. Richlen, Ph.D. of Woods Hole Oceanographic Institution for producing Save Our Ponds and supporting the Cyanobacteria Workshop, Haley Miller U.S. Environmental Protection Agency, and Jasper Sha of Massachusetts Department of Environmental Protection for their review and approval of our Cape Cod Cyanobacteria Monitoring Program (CCCMP) Quality Assurance Project Plan (QAPP) 2024-2029. Amy Usowski, Harwich Conservation Agent, Carrie Schoener, Health Department Director, and Stephanie Ridenour, Director of Natural Resources, for addressing cyanobacteria concerns in Harwich on behalf of the town. Barnstable County Lab for volunteering to sample in the town of Harwich. APCC also thanks our 2025 cyanobacteria interns, including Emma Acri, Sullivan Gaffney, Nora Bowie, Tina Hennig and Sophie Corsaro.

8. REFERENCES

Association to Preserve Cape Cod (APCC) webpage on Cyanobacteria Monitoring Program.
<https://apcc.org/our-work/science/community-science/cyanobacteria/>

Centers for Disease Control (CDC). Facts about Harmful Algal Blooms for Health Care Professionals. <https://stacks.cdc.gov/view/cdc/36648>

Centers for Disease Control and Prevention (CDC). Harmful Algal Bloom (HAB)-Associated Illness. https://www.cdc.gov/harmful-algal-blooms/about/?CDC_AAref_Val=https://www.cdc.gov/habs/general.html

Centers for Disease Control (CDC). Physician Reference card for cyanobacteria.
https://www.cdc.gov/harmful-algal-blooms/media/pdfs/habsphysician_card.pdf

Commonwealth of Massachusetts webpage on cyanobacteria.
<https://www.mass.gov/guides/cyanobacterial-harmful-algal-blooms-cyanohabs-water>

Cyanobacteria Monitoring Collaborative Program (CMC). 2017. Quality Assurance Program Plan (QAPP) for the Cyanobacteria Monitoring Collaborative Program. Rev: 1, June 3, 2021. https://cyanos.org/wp-content/uploads/2024/01/cmc-qapp_2023-07-31.pdf

Environmental Protection Agency (EPA), webpage on Harmful Algal Blooms. https://19january2021snapshot.epa.gov/nutrientpollution/harmful-algal-blooms_.html

EPA and nutrient pollution EPA website on “Monitoring and Responding to Cyanobacteria and Cyanotoxins in Recreational Waters.” <https://www.epa.gov/habs/visually-identifying-signs-cyanobacterial-bloom>

EPA recreational waters EPA Office of Ground Water and Drinking Water webpage. Managing Cyanotoxins in Public Drinking Water Systems. <https://www.epa.gov/ground-water-and-drinking-water/managing-cyanotoxins-public-drinking-water-systems>

Leland, N.J. 2018. Fundamentals of Cyano-casting: cost-effective monitoring techniques for cyanobacteria surface blooms and cyanotoxin levels. In cooperation with UNH Center for Freshwater Biology. http://lim-tex.com/wp-content/uploads/2018/05/CyanoCasting_Handbook_v18.pdf

Leland, N.J. and Haney, J.F. 2018. Alternative Methods for Analysis of Cyanobacterial Populations in Drinking Water Supplies: Fluorometric and Toxicological Applications Using Phycocyanin. Journal of Water Resource and Protection, 10, 740-761. <https://www.scirp.org/journal/PaperInformation?paperID=86671>

Leland, N.J., Haney, J.F., Conte, K., Malkus-Benjamin, K. and Horsley, B. 2019. Evaluation of Size Structure in Freshwater Cyanobacterial Populations: Methods to Quantify Risk Associated with Changes in Biomass and Microcystin Concentrations. Journal of Water Resource and Protection, 11, 810-829. <https://www.scirp.org/journal/paperinformation?paperid=93424>

Leland, N.J., R. A. Landon, and J.F. Haney. September 2020. Trophic interactions between anadromous juvenile Alewife (*Alosa pseudoharengus*) and cyanobacterial populations in a shallow mesotrophic pond. Natural Resources, 2020, 11, 394-419. <https://m.scirp.org/papers/102960>

Local Public Health Institute of Massachusetts. “Key Steps to Collect a Beach Sample.” <https://www.mass.gov/doc/beach-testing-guidelines-for-boards-of-health-and-beach-operators/download>

Massachusetts Department of Public Health (MDPH) website on “Guidelines for cyanobacteria at recreational freshwater locations.” <https://www.mass.gov/info-details/guidelines-for-cyanobacteria-at-recreational-freshwater-locations>

New England Interstate Water Pollution Control Commission (NEIWPCC), webpage on Harmful Algal Blooms. <https://neiwpcc.org/our-programs/wetlands-aquatic-species/habs/>

New Hampshire state issues cyanobacteria advisories and alerts.

<https://www.des.nh.gov/water/healthy-swimming/healthy-swimming-mapper>

New York State Department of Health, Harmful algal bloom program.

<https://www.health.ny.gov/environmental/water/drinking/bluegreenalgae/>

Paerl, H.W., Havens, K.E., Hall, N.S., Otten, T.G., Zhu, M., Xu, H., Zhu, G., and Qin, B. 2019. Mitigating a global expansion of toxic cyanobacterial blooms: confounding effects and challenges posed by climate change. *Marine & Freshwater Research*. Published online March 26, 2019. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7334832/pdf/nihms-1603384.pdf>

Rhode Island Department of Health website on harmful algal blooms.

<https://dem.ri.gov/environmental-protection-bureau/water-resources/research-monitoring/cyanobacteria-blue-green-algae>

State of the Waters: Cape Cod 2020. Action Plan for Homeowners/Business Owners.

<https://capecodwaters.org/action-plan/#ponds-hom>

Town of Barnstable Health Division. Beach Status and Water Quality.

<https://barnstablewaterresources.com/category/water-quality/beach-water-quality-updates/>

U.S. EPA (United States Environmental Protection Agency) (2021). Surface Water Sampling.

https://19january2021snapshot.epa.gov/sites/static/files/2017-07/documents/surface_water_sampling201_af.r4.pdf

U.S. EPA (United States Environmental Protection Agency) (2019). Recommended Human Health Recreational Ambient Water Quality Criteria or Swimming Advisories for Microcystins and Cylindrospermopsin. EPA-822-R-19-001 (PDF).

<https://www.epa.gov/sites/default/files/2019-05/documents/hh-rec-criteria-habs-document-2019.pdf>

U.S. EPA (Environmental Protection Agency). 2016. Method 546: Determination of Total Microcystins and Nodularins in Drinking Water and Ambient Water by Adda Enzyme-Linked Immunosorbent Assay. <https://www.epa.gov/sites/default/files/2016-09/documents/method-546-determination-total-microcystins-nodularins-drinking-water-ambient-water-adda-enzyme-linked-immunosorbent-assay.pdf>

World Health Organization. 1999. Toxic cyanobacteria in water: a guide to their public health consequences, monitoring and management. https://cdn.who.int/media/docs/default-source/wash-documents/water-safety-and-quality/toxic-cyanobacteria---1st-ed.pdf?sfvrsn=338a8c22_1&download=true

WHO (World Health Organization) (2003). Guidelines for Safe Recreational Water Environments, Volume 1: Coastal and Fresh Waters. (PDF)

<https://iris.who.int/bitstream/handle/10665/42591/9241545801.pdf?sequence=1>

9. APPENDIX 1-4

Appendix 1. Resources on Cyanobacteria

Harmful cyanobacteria blooms in freshwater bodies are the subject of numerous reports published by scientists, state and federal agencies, and organizations, some of which are listed here:

- The World Health Organization recognized the public health consequences of cyanobacteria in water in 1999 (https://cdn.who.int/media/docs/default-source/wash-documents/water-safety-and-quality/toxic-cyanobacteria---1st-ed.pdf?sfvrsn=338a8c22_1&download=true).
- The Centers for Disease Control (CDC) call cyanotoxins “among the most powerful natural poisons known” (<https://stacks.cdc.gov/view/cdc/36648>). The https://www.cdc.gov/harmful-algal-blooms/media/pdfs/habsphysician_card.pdf states that swallowing water containing cyanobacteria can damage the central nervous system, liver or kidneys; skin contact can cause allergic dermatitis and conjunctivitis; and inhalation of aerosols containing cyanobacteria or their toxins can cause wheezing, coughing, chest tightness, and shortness of breath.
- New England Interstate Water Pollution Control Commission (<https://neiwpcc.org/our-programs/wetlands-aquatic-species/habs/>) is an interstate commission that helps the states of the Northeast preserve and advance water quality. NEIWPCC’s webpage states that “the frequency of HAB occurrence is on the rise and cyanobacteria toxicity has been associated with human health impacts including skin rashes, gastrointestinal and respiratory disease, and liver damage. Effects can be even more pronounced (potentially even fatal) in animals ranging from cattle to dogs. HABs have direct implications to the use of recreational waterbodies for contact recreation, the susceptibility of public water supplies to toxins, and the overall degradation of our aquatic resources.”
- U.S. Environmental Protection Agency (EPA):
 - “Monitoring and Responding to Cyanobacteria and Cyanotoxins in Recreational Waters.” (<https://www.epa.gov/habs/visually-identifying-signs-cyanobacterial-bloom>)
 - EPA Office of Ground Water and Drinking Water webpage. Managing Cyanotoxins in Public Drinking Water Systems. (<https://www.epa.gov/sites/default/files/2016-09/documents/method-546-determination-total-microcystins-nodularins-drinking-water-ambient-water-adda-enzyme-linked-immunosorbent-assay.pdf>)
 - EPA webpage on nutrient pollution and HABs. (<https://www.epa.gov/habs/visually-identifying-signs-cyanobacterial-bloom>)
 - EPA webpage on Cyanobacteria HABs. (<https://www.epa.gov/habs>)
- State agencies, including New York (<https://www.health.ny.gov/environmental/water/drinking/bluegreenalgae/>), Rhode Island (<https://dem.ri.gov/environmental-protection-bureau/water-resources/research-monitoring/cyanobacteria-blue-green-algae>), and New Hampshire (<https://www.des.nh.gov/water/healthy-swimming/healthy-swimming-mapper>) have

cyanobacteria monitoring programs and provide guidance concerning public health and environmental risks posed by cyanobacteria.

- Commonwealth of Massachusetts:
 - Cyanobacteria webpage: (<https://www.mass.gov/guides/cyanobacterial-harmful-algal-blooms-cyanohabs-water>)
 - Massachusetts Department of Public Health (MDPH) website on “Guidelines for cyanobacteria in freshwater recreational water bodies.” (<https://www.mass.gov/info-details/guidelines-for-cyanobacteria-at-recreational-freshwater-locations>)

Appendix 2. APCC's Cyanobacteria Risk Categories.

Risk Category Descriptions

Acceptable

Definition: No concerning cyanobacteria results at the time and place of sampling. To the best of APCC's knowledge and monitoring results, toxin levels are below State standards for recreational usage of the pond with respect to cyanobacteria and toxins.

Map color is blue.

Formerly the Low Warning Tier.

Recommended Sampling Frequency: Biweekly.

Recommended Action: None.

Potential for Concern

Definition: Monitoring results or the presence of cyanobacteria scum at the time and place of sampling indicate a potential for increased risk for exposure to cyanobacteria toxins. While these conditions may pose lower health risks to adults, risks are higher for children or pets based on lower body mass, particularly if contaminated water is incidentally ingested. Children may inadvertently consume pond water while swimming and pet exposure can result from drinking or ingesting pond water or from grooming after swimming.

Map color is yellow.

Map color yellow with crosshatching indicates a municipal pet advisory has been issued.

Formerly the Moderate Warning Tier

Sampling Frequency: Weekly, unless APCC is notified to hold off on weekly sampling by the client.

Recommended Action:

1. APCC will screen a GRAB sample for microcystin using toxin strip test if microscopy determines the sample is *Microcystis* dominant. If the screening result is equal to or higher than 4ppb APCC will provide a GRAB sample for toxin analysis to the Barnstable County Water Quality Lab for confirmatory testing.
2. A "Pet Advisory" or similar advisory may be posted in accordance with each town's policies and procedures until the pond returns to Acceptable status.
3. Sampling is increased to weekly until all results are once again in the "Acceptable" category, unless APCC is notified to hold off on weekly sampling by the client.

Use Restriction Warranted

Definition: Monitoring results at the time and place of sampling indicate the pond is unsafe for recreation by humans and pets based on the presence of microcystin at or above state standards (8 ppb microcystin) OR based on determination by a municipal health agent(s) resulting in a closure for any other reason related to cyanobacteria. Recreational risk to adults is moderate following exposure. Recreational risks are especially high for children and pets following exposure through accidental ingestion of contaminated water. Children may inadvertently consume pond water while swimming and pet exposure can result from ingestion or directly drinking pond water or from grooming after swimming. Due to lower body masses, children and pets are more susceptible to cyanobacteria risks than adults.

Map color is red.

Map color red with crosshatching indicates a municipal advisory has been issued.

Formerly the High Warning Tier.

Sampling Frequency: Weekly, unless APCC is notified to hold off on weekly sampling by the client.

Recommended Action:

1. APCC will screen a GRAB sample for microcystin using toxin strip test if microscopy determines the sample is *Microcystis* dominant. If the screening result is equal to or higher than 4ppb APCC will provide a GRAB sample for toxin analysis to the Barnstable County Water Quality Lab for confirmatory testing.
2. The town should post a recreational advisory or similar advisory according to municipal policies and procedures and otherwise notify the public to avoid contact and exposure until the pond meets criteria to be reopened or the advisory is lifted by the local health agent.
3. Sampling will be conducted weekly (unless APCC is notified by client to hold off on weekly sampling) and the pond will remain in the “Use Restriction Warranted” category until there are two consecutive “Acceptable” sampling events according to APCC’s 2025 Risk Tier Table.

APCC 2025 Cyanobacteria Risk Tiers

Acceptable

- A cyanobacteria scum was not detected, and the Bloom Forming Colony sample had a phycocyanin measurement <500ug/L.
- A cyanobacteria scum was detected but was determined to be visually insignificant and the Bloom Forming Colony sample had a phycocyanin measurement <100ug/L.

Potential for Concern

- A cyanobacteria scum was not detected but the Bloom Forming Colony sample had a phycocyanin measurement ≥500ug/L.
- A cyanobacteria scum was detected and determined to be visually significant and/or the Bloom Forming Colony sample had a phycocyanin measurement ≥100ug/L.

Stripes are added on the map if the town posts a warning that is not an official Public Health Advisory.*

Use Restriction Warranted

- The town posts a Public Health Advisory.
- Microcystin test measures ≥8 ppb (MassDPH guidance).
- Once a pond is categorized as Use Restriction Warranted it will remain in this category for two consecutive Acceptable sampling events (MassDPH guidance).

Stripes are added on the map if the town posts a Public Health Advisory and are removed once the town removes the Public Health Advisory. **If and when the town informs APCC of their action*

Contact: Julie Hambrook, APCC Cyanobacteria Monitoring Program Email: jhambrook@apcc.org

Appendix 3. Bucks Pond, Cornelius Pond, Hawksnest Pond, Hinkley Pond, Josephs Pond, Sand Lake, Sand Pond, Skinequit Pond, and West Reservoir 2021-2025 Cyanobacteria Risk Comparison.

As of 2025, APCC has completed five seasons of cyanobacteria monitoring for the town of Harwich. The tables below detail APCC’s communication of cyanobacteria risk for Bucks Pond, Cornelius Pond, Hawksnest Pond, Hinkley Pond, Josephs Pond, Sand Lake, Sand Pond, Skinequit Pond, and West Reservoir each season. Red indicates a “Use Restriction Warranted” or “High Warning Tier” designation and yellow indicates a “Potential for Concern” or “Moderate Warning Tier” designation, and blue indicates an “Acceptable” or “Low Warning Tier” designation. The coloring indicates the worst-case from any location in the pond. Even if a sampling event took place outside of the pictured time window, the worst risk tier displayed on the map during that time window was used to create the tables. See previous APCC reports for the town of Harwich for more information on findings and risk communication in these sampling seasons.

Bucks Pond 2021-2025 Cyanobacteria Risk Comparison												
	June		July		August		September		October		November	
Year	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-31st	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-30th
2021	Blue	Blue	Blue	Blue	Blue	Blue	Blue	Blue	Blue			
2022	Blue	Blue	Yellow	Yellow	Blue	Blue						
2023	Blue	Blue	Blue	Blue	Blue	Blue						
2024	Blue	Blue	Yellow	Blue	Blue							
Note that APCC updated the risk assessment framework in 2025.												
2025	Blue	Yellow	Yellow	Blue	Blue	Blue	Blue	Blue	Blue	Blue		

Cornelius Pond 2024-2025 Cyanobacteria Risk Comparison												
	June		July		August		September		October		November	
Year	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-31st	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-30th
2024	Blue	Blue	Blue	Blue	Blue							
Note that APCC updated the risk assessment framework in 2025.												
2025	Blue	Blue	Blue	Blue	Blue							

Hawksnest Pond 2024-2025 Cyanobacteria Risk Comparison												
	June		July		August		September		October		November	
Year	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-31st	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-30th
2024	Blue	Blue	Blue	Blue	Blue							
Note that APCC updated the risk assessment framework in 2025.												
2025	Blue	Blue	Blue									

Hinckleys Pond 2021-2025 Cyanobacteria Risk Comparison												
	June		July		August		September		October		November	
Year	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-31st	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-30th
2021												
2022												
2023												
2024												
Note that APCC updated the risk assessment framework in 2025.												
2025												

Josephs Pond 2021-2025 Cyanobacteria Risk Comparison												
	June		July		August		September		October		November	
Year	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-31st	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-30th
2021												
2022												
2023												
2024												
Note that APCC updated the risk assessment framework in 2025.												
2025												

Sand Lake 2024-2025 Cyanobacteria Risk Comparison												
	June		July		August		September		October		November	
Year	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-31st	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-30th
2024												
Note that APCC updated the risk assessment framework in 2025.												
2025												

Sand Pond 2025 Cyanobacteria Risk Comparison												
	June		July		August		September		October		November	
Year	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-31st	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-30th
2025												

Skinequit Pond 2021-2025 Cyanobacteria Risk Comparison												
	June		July		August		September		October		November	
Year	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-31st	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-30th
2021												
2022												
2023												
2024												
Note that APCC updated the risk assessment framework in 2025.												
2025												

West Reservoir 2021-2025 Cyanobacteria Risk Comparison												
	June		July		August		September		October		November	
Year	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-31st	1st-15th	16th-30th	1st-15th	16th-31st	1st-15th	16th-30th
2021												
2022												
2023												
2024												
Note that APCC updated the risk assessment framework in 2025.												
2025												

Appendix 4. Sampling Locations

- Bucks Pond – Pleasant Park Circle (Private).
GPS Coordinates: 41.70010253, -70.04329141
- Cornelius Pond – Residence off Eldridge Pond Road (Private).
GPS Coordinates: 41.70548951, -70.06050064
- Hawksnest Pond – Parking off Nathan Walker Road.
GPS Coordinates: 41.7137019, -70.04488397
- Hinckleys Pond – Off Cape Cod Rail Trail.
GPS Coordinates: 41.71306274, -70.0898452
- Josephs Pond – Neighborhood beach off Vacation Lane (Private).
GPS Coordinates: 41.70079314, -70.04941511
- Sand Lake – Path to private beach off Rainbow Way.
GPS Coordinates: 41.70045968, -70.03663149
- Sand Pond –
GPS Coordinates: 41.68925927, -70.11212731
- Skinequit Pond – Neighborhood landing off Ocean Street (Private).
GPS Coordinates: 41.67375157, -70.04290988
- West Reservoir – Bells Neck Road off Great Western Road.
GPS Coordinates: 41.68430085, -70.11867274